

CO Emission in Powerful Radio Galaxies at Low and High Redshift

A. S. Evans

California Institute of Technology 105-24, Pasadena, CA 91125 U.S.A.

Abstract. CO observations of $0 < z < 4$ radio galaxies are presented. The L_{CO} upper limits for the high-redshift powerful radio galaxies (HzPRGs: $z > 1$, $P_{408\text{MHz}} > 10^{27}$ Watts Hz^{-1}) are consistent with no evolution in molecular gas (H_2) mass with redshift and/or radio power. Of the low-redshift powerful radio galaxies (LzPRGs: $z < 0.2$, $P_{408\text{MHz}} > 10^{23.5}$ W Hz^{-1}) observed, only one F-R II galaxy has been detected in CO, whereas $\sim 50\%$ of the radio-compact (FC) and F-R I galaxies have been detected. The CO and imaging data to date imply that either F-R II galaxies result from mergers of gas-poor galaxies relative to FC/F-R I galaxies, or that some FC/F-R I galaxies evolve into F-R II galaxies. In either scenario, the radio activity in powerful radio galaxies is most likely triggered by a merger event, and H_2 may be the fuel for the active galactic nucleus (AGN) in the early stages of the merger.

1. Introduction

Advances in millimeter receiver technology and observing techniques, as well as the availability of moderate-size millimeter telescopes and arrays, have made recent CO observations of distant extragalactic sources possible. During the mid-1980s, CO surveys of galaxies selected by their *IRAS* fluxes and colors resulted in CO detections of a host of $z < 0.2$ Seyfert, HII region-like, and LINER galaxies, as well as several QSOs and radio galaxies (e.g., Sanders, Scoville, & Soifer 1991). Their inferred H_2 masses are $1 \times 10^9 - 6 \times 10^{10} M_{\odot}$. For galaxies believed to have AGNs, H_2 provides a possible source of fuel for nuclear activity, as well as for vigorous, circumnuclear star formation. The detection of CO emission in the $z = 2.3$ *IRAS* galaxy FSC 10214+4724 (Brown & Vanden Bout 1991, 1992) revolutionized the field, both because of its cosmological implications and because it proved that it was possible, with modern technology, to detect star-forming gas out to redshifts of a few. Since then, much effort has gone into searching for CO in a wide variety of high-redshift objects.

At the time FSC 10214+4724 was detected in CO, the only sample of galaxies known at comparable redshifts were radio galaxies. HzPRGs were well suited as initial targets for CO surveys, primarily because (i) several LzPRGs had been detected in CO (Phillips et al. 1987; Mirabel, Sanders, & Kazès 1989; Mazzarella et al. 1993), (ii) radio galaxies are observed to span a wide redshift range ($0 < z < 4$) and (iii) their redshifts are known to the accuracy required by the 512 MHz bandwidths of millimeter detectors. Although the




Figure 1. L'_{CO} and $M(\text{H}_2)$ vs. z (Evans et al. 1996). $\downarrow$ denotes 3σ upper limits. $H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$ is assumed.

number of LzPRGs observed in CO at that time was small, further evidence that radio galaxies as a class may be rich in H_2 came from the observations that a significant fraction (20–30%) of LzPRGs were known to be infrared luminous (Golombek, Miley, & Neugebauer 1988). Some have infrared excesses consistent with thermal emission from dust, and many LzPRGs possess morphological peculiarities commonly associated with collisions/mergers of galaxy pairs (Heckman et al. 1986). Multi-component structures and/or nearby companions are also observed in $z \sim 1$ HzPRGs (Spinrad & Djorgovski 1984; Lilly & Longair 1984; Djorgovski et al. 1987), but the *IRAS* flux upper limits yield very little information on the shape and luminosity of the far-infrared emission in these galaxies (Evans et al. 1996). The blue colors of many HzPRGs are often cited as additional, though less compelling, evidence that they may be H_2 rich; the colors may also be due to scattered AGN light.

In this article, the results of the CO observations of HzPRGs (e.g., Evans et al. 1996; Downes et al. 1996) will first be summarized, followed by a discussion of a complete 60 and 100 μm flux-limited sample of LzPRGs (e.g. Evans et al. 1998a). The latter discussion will aim at clarifying the results obtained from the high-redshift observations. With the exception of Figure 1, $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $q_0 = 0.5$, and $\alpha = 4 \text{ M}_\odot (\text{K km s}^{-1} \text{ pc}^2)^{-1}$ are assumed throughout.

2. The High-Redshift Radio Galaxy Survey

The primary goals of any CO survey of high-redshift galaxies are to estimate the H_2 masses and to determine if the H_2 content evolves as a function of redshift. If the radio power of radio galaxies is correlated with the amount of H_2 , then HzPRGs, which are preselected by their high radio luminosity, may have rich circumnuclear ISMs that serve as fuel for the AGN. Thus, a sample of 15 HzPRGs spanning the redshift range $1 < z < 4$ were selected. For each galaxy, observations were done by tuning to $J > 1$ rotational CO transitions redshifted




Figure 2. Examples of CO emission lines in LzPRGs (Evans et al. 1998a).

to frequencies accessible by the 3mm receivers on the NRAO 12m and IRAM 30m Telescopes. Eleven HzPRGs were eventually observed at one or both telescopes, with integration times of up to 10 hours at the 30m Telescope, and up to 50 hours at the 12m Telescope.

None of the eleven galaxies observed were detected (Figure 1). However, assuming a line width of $v_{\text{FWHM}} \sim 250 \text{ km s}^{-1}$ (the average v_{FWHM} observed for a complete sample of ultraluminous infrared galaxies, ULIGs: Sanders, Scoville, & Soifer 1991), the 3σ upper limits for H_2 mass are in the range $3.5\text{--}20 \times 10^{10} M_{\odot}$ ($H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$). Further, the four HzPRGs with the most sensitive CO upper limits would appear to have less H_2 than the most H_2 -rich galaxies in the local universe. This would seem to imply that the H_2 content of radio galaxies does not evolve as a function of redshift or radio power.

Is it the case that, while radio-weak ULIGs show evidence of evolution of H_2 mass with redshift, radio-loud infrared galaxies, such as radio galaxies, do not? Indeed, the recent tentative detection of CO in 53W002 ($\sim 7 \times 10^{10} M_{\odot}$ of H_2 : Scoville et al. 1997), a relatively radio-faint HzPRG in comparison with the 11 HzPRGs discussed above, may be evidence of an anticorrelation between H_2 mass and radio power. In actuality, there is very little evidence for evolution of H_2 mass in any class of galaxies as a function of redshift. IRAS 10214+4724 is now known to be gravitationally lensed (e.g. Eisenhardt et al. 1996); its inferred intrinsic H_2 mass is $\sim 10^{10} M_{\odot}$, a fairly average value for an ULIG. In 1994, a $z = 2.6$ gravitationally-lensed QSO (H1413+117) was detected (Barvainis et al. 1994), with an intrinsic luminosity also believed to be average for ULIGs at low redshift. Two other ULIGs (IR 09104+4109 at $z = 0.4$ and IR 15307+3252 at $z = 0.9$) and one high-redshift *IRAS*-luminous QSO (PG 1634+706 at $z = 1.3$) have been searched for CO, but the 3σ upper limits of L'_{CO} are less than that of the most gas-rich galaxies observed locally (Evans et al. 1998b). Finally, the $z = 4.7$ QSO BR 1202-07 has recently been detected and confirmed, and its inferred H_2 mass is $\sim 1 \times 10^{11} M_{\odot}$ (Omont et al. 1996; Ohta et al. 1996). This may be the first indication of evolution in H_2 mass with redshift, however, the

Figure 3. The F-R II galaxy 3C 293 (Evans et al. 1999). The scale is 860 pc/". The radio continuum map is from Leahy, Pooley, & Riley (1986).

H₂ mass of BR 1202-07 is only twice that observed in the most H₂-rich, local galaxies. There is also evidence for shear due to gravitational lensing in the vicinity of the QSO, which may indicate that its intrinsic L'_{CO} is much lower than observed (Omont et al. 1996).

If, as is commonly believed, radio galaxies are the progenitors of some giant elliptical galaxies, there are four possible reasons why CO emission has not been unambiguously detected in any HzPRGs, and why evolution of H₂ mass with redshift is not seen in other classes of galaxies: First, the redshift at which the bulk of stars in present day galaxies formed may be greater than four. Thus, most of the H₂ may have been consumed by starbursts and the AGN by the redshift most of these galaxies are located. Second, because merger events must have been more common in the past, most of the H₂ that eventually formed the bulk of the stars in giant elliptical galaxies may be in many smaller, star-

forming galaxies. These individual galaxies may consume most of their H_2 before merging. Third, the duration that galaxies can contain in excess of $4 - 6 \times 10^{10} M_\odot$ of H_2 may be very brief. Thus, the probability of observing such a gas-rich galaxy may be exceedingly low. Fourth, the universe may not have closure density, thus the scale of the universe may be larger than assumed for the above H_2 mass calculations. Even given this possibility, the H_2 mass upper limits of the four best studied HzPRGs are substantially less than the total stellar mass of a giant elliptical galaxy.

3. The Low-Redshift Radio Galaxy Survey

From the high-redshift observations alone, it was not clear what role H_2 may play in the radio activity seen in these galaxies. In addition, the number of radio galaxies observed prior to the HzPRG CO survey was very small, thus it was not clear if inaccurate generalizations based on observations of a few gas-rich LzPRGs were being made for radio galaxies as a whole. Of the several LzPRGs that had been observed, none of those detected had been F-R II (i.e., edge-brightened radio morphology, typically observed to have $P_{408\text{MHz}} > 10^{25.3} \text{ W Hz}^{-1}$; Fanaroff & Riley 1974) galaxies like the HzPRGs - all had been FC or F-R I (i.e., edge-darkened radio morphologies, typically with $P_{408\text{MHz}} < 10^{25.3} \text{ W Hz}^{-1}$) galaxies.

Over the last few years, considerable time has been devoted to investigating the H_2 masses and morphologies, as well as the $2.2\mu\text{m}$ morphologies of a complete sample of 35 LzPRGs with *IRAS* fluxes at $60\mu\text{m}$ and/or $100\mu\text{m} > 0.30 \text{ Jy}$ and $\delta > -40^\circ$ (Evans et al. 1998a). Examples of some of the CO emission lines detected are shown in Figure 2. The single-dish observations are being obtained using the NRAO 12m Telescope, with follow-up interferometry of detected sources obtained using the Owens Valleys Millimeter Array. The $2.2\mu\text{m}$ imaging, obtained to look for evidence of isophotal irregularities, imbedded multiple nuclei and bridges hidden by dust at optical wavelengths, was done using the UH 2.2m Telescope. The results from this survey are that only 40% of the F-R II galaxies show evidence of disturbed, high surface brightness features and multiple nuclei or close companions and that, aside from a CO detection in 3C 293 ($z \sim 0.04$; Evans et al. 1999, Figure 2 and 3), none of the F-R II sources have confirmed CO detections. Cygnus A, the prototypical F-R II radio galaxy, is among the galaxies that were not detected; although it has an $L'_{\text{IR}} \sim 4.5 \times 10^{11} L_\odot$, it has less than one-third the H_2 mass of the Milky Way. The $2.2\mu\text{m}$ image (not shown) supports this result - the host galaxy looks like an undisturbed elliptical galaxy. Further, Cygnus A and many other LzPRGs have infrared colors suggestive of warm dust temperatures, indicating that most of the infrared luminosity may be coming from a small amount of dust heated by the AGN (i.e., $L_{\text{IR}} \propto M_{\text{dust}} T_{\text{dust}}^5$). In contrast, 86% of the FC and F-R I sources show evidence of disturbed, high surface brightness features and multiple nuclei or close companions, and 50% have been detected in CO. Such a dramatic contrast could simply suggest that the progenitors of F-R II galaxies are gas-poor relative to the progenitors of FC and F-R I sources, or that some FC and F-R I sources evolve into F-R II sources after the H_2 has been consumed. The fundamental difference between F-R I and F-R II sources may be the efficiency to which the

jets propagate through the ISM of the host galaxy. Given this, the lack of CO detections in HzPRGs is not surprising.

If the apparent deficiency in the H_2 masses of F-R II relative to FC/F-R I galaxies is due to time evolution, galaxies such as 3C 293 may be transitional sources. The galaxy 3C 293 (Figure 3) has a moderately high infrared luminosity, a distorted disk-like infrared morphology with a companion, and several times the H_2 mass of the Milky Way. The CO emission is extended along the axis of the $2.2\ \mu\text{m}$ disk and perpendicular to the radio jet, possibly because the jets have escaped through an axis unobscured by the putative nuclear torus. The CO emission line (Figure 2), has an absorption feature approximately $150\ \text{km s}^{-1}$ wide. If the width of the line is due solely to rotation, the absorption may occur from gas $\sim 50\ \text{pc}$ from the AGN, approximately the radius of the dust torus observed in HST images of the radio galaxy NGC 4261 (Jaffe et al. 1996).

Acknowledgments. It is a pleasure to thank my collaborators D. Sanders, J. Mazzarella, P. Solomon, D. Downes, S. Simon, J. Surace, F. Mirabel, and C. Kramer. I also am grateful to the staffs of the NRAO 12m, IRAM 30m, and UH 2.2m Telescopes, and the staff and postdocs of the Owens Valley Millimeter Array for their assistance during and after the observations presented here.

References

- Barvainis, R. et al. 1994, *Nature*, 371, 586
Djorgovski, S. et al. 1987, *AJ*, 93, 1307
Downes, D., Solomon, P., Sanders, D. B., & Evans, A. S. 1996, *A&A*, 313, 91
Eisenhardt, P. R. et al. 1996, *ApJ*, 461, 72
Evans, A. S. et al. 1996, *ApJ*, 457, 658
Evans, A. S. et al. 1998a, *ApJ*, in prep.
Evans, A. S. et al. 1998b, *ApJ*, 506, 205
Evans, A. S. et al. 1999, *ApJ*, in press
Fanaroff, B. L. & Riley, F. M. 1974, *MNRAS*, 167, 31P
Golombek, D., Miley, G. K., & Neugebauer, G. 1988, *AJ*, 95, 26
Heckman, T. M. et al. 1986, *ApJ*, 311, 526
Jaffe, W. et al. 1996, *ApJ*, 460, 214
Leahy, J. P., Pooley, G. G., & Riley, J. M. 1986, *MNRAS*, 222, 753
Lilly, S. & Longair, M. 1984, *MNRAS*, 211, 833
Mazzarella, J. M. et al. 1993, *ApJ*, 409, 170
Mirabel, I. F., Sanders, D. B., & Kazès, I. 1989, *ApJ*, 340, L9
Ohta, K. et al. 1996, *Nature*, 382, 426
Omont et al. 1996, *Nature*, 382, 428
Phillips, T. G. et al. 1987, *ApJ*, 322, L73
Sanders, D. B., Scoville, N. Z., & Soifer, B. T. 1991, *ApJ*, 370, 158
Scoville, N. Z. et al. 1997, *ApJ*, 485, L21
Spinrad, H., & Djorgovski, S. G. 1984, *ApJ*, 280, L9

This figure "evansa_fig3.gif" is available in "gif" format from:

<http://arXiv.org/ps/astro-ph/9810163v1>